

Table 1: Direct experimental measurements of $\phi_s^{c\bar{c}s}$, $\Delta\Gamma_s$ and Γ_s using $B_s^0 \rightarrow J/\psi \phi$, $J/\psi K^+K^-$, $J/\psi \pi^+\pi^-$ and $D_s^+D_s^-$ decays. Only the solution with $\Delta\Gamma_s > 0$ is shown, since the two-fold ambiguity has been resolved in Ref. [1]. The first error is due to statistics, the second one to systematics. The last line gives our average.

Exp.	Mode	Dataset	$\phi_s^{c\bar{c}s}$	$\Delta\Gamma_s$ (ps $^{-1}$)	Ref.
CDF	$J/\psi \phi$	9.6 fb $^{-1}$	[−0.60, +0.12], 68% CL	+0.068 ± 0.026 ± 0.009	[2]
D0	$J/\psi \phi$	8.0 fb $^{-1}$	−0.55 $^{+0.38}_{-0.36}$	+0.163 $^{+0.065}_{-0.064}$	[3]
ATLAS	$J/\psi \phi$	4.9 fb $^{-1}$	+0.12 ± 0.25 ± 0.05	+0.053 ± 0.021 ± 0.010	[4]
ATLAS	$J/\psi \phi$	14.3 fb $^{-1}$	−0.119 ± 0.088 ± 0.036	+0.096 ± 0.013 ± 0.007	[5] ^p
ATLAS	above 2 combined		−0.094 ± 0.083 ± 0.033	+0.082 ± 0.011 ± 0.007	[5] ^p
CMS	$J/\psi \phi$	20 fb $^{-1}$	−0.075 ± 0.097 ± 0.031	+0.095 ± 0.013 ± 0.007	[6]
LHCb	$J/\psi K^+K^-$	3.0 fb $^{-1}$	−0.058 ± 0.049 ± 0.006	+0.0805 ± 0.0091 ± 0.0033	[7]
LHCb	$J/\psi \pi^+\pi^-$	3.0 fb $^{-1}$	+0.070 ± 0.068 ± 0.008	—	[8]
LHCb	above 2 combined		−0.010 ± 0.039(tot)	—	[7]
LHCb	$D_s^+D_s^-$	3.0 fb $^{-1}$	+0.02 ± 0.17 ± 0.02	—	[9]
All combined			−0.034 ± 0.033	+0.084 ± 0.007	

^p Preliminary.

References

- [1] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. Lett. **108**, 241801 (2012), arXiv:1202.4717 [hep-ex].
- [2] T. Aaltonen *et al.* (CDF collaboration), Phys. Rev. Lett. **109**, 171802 (2012), arXiv:1208.2967 [hep-ex]; this replaces T. Aaltonen *et al.* (CDF collaboration), Phys. Rev. **D85**, 072002 (2012), arXiv:1112.1726 [hep-ex]; as well as T. Aaltonen *et al.* (CDF collaboration), Phys. Rev. Lett. **100**, 161802 (2008), arXiv:0712.2397 [hep-ex]; and T. Aaltonen *et al.* (CDF collaboration), Phys. Rev. Lett. **100**, 121803 (2008), arXiv:0712.2348 [hep-ex].
- [3] V. M. Abazov *et al.* (D0 collaboration), Phys. Rev. **D85**, 032006 (2012), arXiv:1109.3166 [hep-ex]; this replaces Ref. [?] and V. M. Abazov *et al.* (D0 collaboration), Phys. Rev. Lett. **101**, 241801 (2008), arXiv:0802.2255 [hep-ex]; as well as V. M. Abazov *et al.* (D0 collaboration), Phys. Rev. Lett. **98**, 121801 (2007), arXiv:hep-ex/0701012.
- [4] G. Aad *et al.* (ATLAS collaboration), Phys. Rev. **D90**, 052007 (2014), arXiv:1407.1796 [hep-ex]; this replaces G. Aad *et al.* (ATLAS collaboration), JHEP **12**, 072 (2012), arXiv:1208.0572 [hep-ex].
- [5] J. Walder (for the ATLAS collaboration), CERN seminar, July 7, 2015, <http://indico.cern.ch/event/388146>.

- [6] V. Khachatryan *et al.* (CMS collaboration), submitted to Phys. Lett. **B** (2015), arXiv:1507.07527 [hep-ex]; this supersedes CMS collaboration, CMS note CMS-PAS-BPH-13-012, 22 July 2014, <https://cds.cern.ch/record/1744869>.
- [7] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. Lett. **114**, 041801 (2015), arXiv:1411.3104 [hep-ex]; this replaces any $B_s^0 \rightarrow J/\psi K^+ K^-$, $B_s^0 \rightarrow J/\psi \phi$ or combined $\phi_s^{c\bar{c}s}$ result from Ref. [10].
- [8] R. Aaij *et al.* (LHCb collaboration), Phys. Lett. **B736**, 186 (2014), arXiv:1405.4140 [hep-ex]; this replaces any $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ or $B_s^0 \rightarrow J/\psi f_0(980)$ result on CP violation from Ref. [10].
- [9] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. Lett. **113**, 211801 (2014), arXiv:1409.4619 [hep-ex].
- [10] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. **D87**, 112010 (2013), arXiv:1304.2600 [hep-ex]; this supersedes the following publications, R. Aaij *et al.* (LHCb collaboration), Phys. Rev. Lett. **108**, 101803 (2012), arXiv:1112.3183 [hep-ex]; R. Aaij *et al.* (LHCb collaboration), Phys. Lett. **B713**, 378–386 (2012), arXiv:1204.5675 [hep-ex]; R. Aaij *et al.* (LHCb collaboration), Phys. Lett. **B707**, 497–505 (2012), arXiv:1112.3056 [hep-ex]; this also replaces the $B_s^0 \rightarrow J/\psi f_0(980)$ result of R. Aaij *et al.* (LHCb collaboration), Phys. Rev. Lett. **109**, 152002 (2012), arXiv:1207.0878 [hep-ex].