List of other measurements that are not included in the tables:

- $B^+ \to K^+ \pi^- \pi^+ \gamma$: LHCb has measured the up-down asymmetries in bins of the $K\pi\pi\gamma$ mass [91].
- In [100], LHCb has also measured the BR of $B^+ \to K^+ e^- e^+$ in the $m^2(\ell\ell)$ bin [1, 6] $(\text{GeV}/c^2)^2$.
- In the $B^+ \to \pi^+ \mu^+ \mu^-$ paper [92], LHCb has also measured the differential BR in bins of $m^2(\ell\ell)$.
- For $B \to K\ell^-\ell^+$, LHCb has measured F_H and A_{FB} in 17 (5) bins of $m^2(\ell\ell)$ for the K^+ (K_s) final state [105]. Belle has measured F_L and A_{FB} in 6 $m^2(\ell\ell)$ bins [64].
- For the $B \to K^* \ell^- \ell^+$ analyses, partial branching fractions and angular observables in bins of $m^2(\ell\ell)$ are also available :
 - $-B^0 \to K^{*0}e^-e^+$: LHCb has measured F_L , $A_T^{(2)}$, A_T^{Im} , A_T^{Re} in the $m^2(\ell\ell)$ bin [0.002, 1.120] $(\text{GeV}/c^2)^2$ [101], and the BR in the dilepton mass region [10,1000] (MeV/c^2) [103].
 - $-B \to K^*\ell^-\ell^+$: Belle has measured F_L , A_{FB} , isospin asymetry in 6 $m^2(\ell\ell)$ bins [41] and P_4' , P_5' , P_6' , P_8' in 4 $m^2(\ell\ell)$ bins [64]. Babar has measured F_L , A_{FB} , P_2 in 5 $m^2(\ell\ell)$ bins [37].
 - $-B^0 \to K^{*0} \mu^- \mu^+$: LHCb has measured F_L , A_{FB} , $S_3 S_9$, $A_3 A_9$, $P_1 P_3$, $P_4' P_8'$ in 8 $m^2(\ell\ell)$ bins [104]. CMS has measured F_L , A_{FB} in 7 $m^2(\ell\ell)$ bins [115]. ATLAS has measured F_L and A_{FB} in 5 $m^2(\ell\ell)$ bins [114] (conference note).