

Heavy FLavor AVeraging group (HFLAV) - May 2018

Compilation of Λ_b^0 Branching Fractions ($\times 10^{-6}$) - UL at 90% CL

Preliminary Updated results not included in PDG Live as of Dec. 31, 2017

RPP#	Mode	PDG2017 Avg.	CDF	LHCb	Our Avg.
12	$\bar{K}^0 p \pi^-$	13.0 ± 4.0		$12.6 \pm 1.9 \pm 0.9 \pm 3.4 \pm 0.5$ § [1]	12.6 ± 4.0
13	$K^0 p K^-$	< 3.5		< 3.5 [1]	< 3.5
33	$p \pi^-$	4.3 ± 0.8	$3.5 \pm 0.6 \pm 0.9$ [2]		3.5 ± 1.1
34	$p K^-$	5.1 ± 0.9	$5.6 \pm 0.8 \pm 1.5$ [2]		5.6 ± 1.7
37	$\Lambda \mu^+ \mu^-$	1.08 ± 0.28	$1.73 \pm 0.42 \pm 0.55$ [3]	$0.96 \pm 0.16 \pm 0.25$ [4]	1.08 ± 0.27
38	$\Lambda \gamma$	< 1300	< 1300 [5]		< 1300
39	$\Lambda \eta$	9^{+7}_{-5}		$9.3^{+7.3}_{-5.3}$ ¶ [6]	$9.3^{+7.3}_{-5.3}$
40	$\Lambda \eta'$	< 3.1		< 3.1 [6]	< 3.1
41	$\Lambda \pi^+ \pi^-$	4.7 ± 1.9		$4.6 \pm 1.2 \pm 1.4 \pm 0.6$ † 2 [7]	4.6 ± 1.9
42	$\Lambda K^+ \pi^-$	5.7 ± 1.3		$5.6 \pm 0.8 \pm 0.8 \pm 0.7$ † 2 [7]	5.6 ± 1.3
43	$\Lambda K^+ K^-$	16.1 ± 2.3		$15.9 \pm 1.2 \pm 1.2 \pm 2.0$ † 2 [7]	15.9 ± 2.6
44	$\Lambda \phi$	2.0 ± 0.5		$5.18 \pm 1.04 \pm 0.35^{+0.67}_{-0.62} \pm 3$ [8]	$5.18^{+1.29}_{-1.26}$
	$\Lambda p \pi^- \mu^+ \mu^-$			$0.069 \pm 0.019 \pm 0.011^{+0.013}_{-0.010}$ † [9]	$0.069^{+0.026}_{-0.024}$
	$\Lambda p \pi^- \pi^+ \pi^-$			$19.0 \pm 0.6 \pm 1.0 \pm 1.6 \pm 0.7$ † [10]	19.0 ± 2.1
	$\Lambda p K^- \pi^+ \pi^-$			$45.5 \pm 0.8 \pm 2.0 \pm 3.9 \pm 1.7$ † [10]	45.5 ± 4.8
	$\Lambda p K^- K^+ \pi^-$			$3.7 \pm 0.3 \pm 0.4 \pm 0.3 \pm 0.1$ † [10]	3.7 ± 0.6
	$\Lambda p K^- K^+ K^-$			$11.4 \pm 0.3 \pm 0.7 \pm 1.0 \pm 0.5$ † [10]	11.4 ± 1.4

Channels with no RPP# are not reported by PDG.

Results for CDF and LHCb are relative BFs converted to absolute BFs.

† Last quoted uncertainty is due to the precision with which the normalization channel branching fraction is known.

‡ Third uncertainty is related to external inputs.

§ Third uncertainty is from the ratio of fragmentation fractions $f_{\Lambda_b^0}/f_d$, and the fourth is due to the uncertainty on $\mathcal{B}(B^0 \rightarrow K^0 \pi^+ \pi^-)$.

¶ Result at 68% CL.

¹ Third uncertainty is from $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ p \pi^-)$, and the fourth is due to the uncertainty on $\mathcal{B}(\Lambda_c^+ \rightarrow p K^- \pi^+)$.

² Normalization taken directly from LHCb paper.

³ Difference w.r.t. PDG value under investigation.

Heavy FLavor AVeraging group (HFLAV) - May 2018

Partial Branching Fractions of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ decays ($\times 10^{-7}$)

Preliminary Updated results not included in PDG Live as of Dec. 31, 2017

Mode	q^2 [GeV^2/c^4] † ‡	PDG2017 Avg.	CDF	LHCb	Our Avg.
$\Lambda \mu^+ \mu^-$	< 2.0	0.71 ± 0.27	$0.15 \pm 2.01 \pm 0.05$ [3]	$0.72^{+0.24}_{-0.22} \pm 0.14$ [11]	$0.71^{+0.27}_{-0.26}$
$\Lambda \mu^+ \mu^-$	[2.0, 4.3]	$0.28^{+0.28}_{-0.21}$	$1.8 \pm 1.7 \pm 0.6$	$0.253^{+0.276}_{-0.207} \pm 0.046$ [11]	$0.281^{+0.286}_{-0.218}$
$\Lambda \mu^+ \mu^-$	[4.3, 8.68]	0.5 ± 0.7	$-0.2 \pm 1.6 \pm 0.1$	$0.66 \pm 0.72 \pm 0.16$ [4]	0.51 ± 0.67
$\Lambda \mu^+ \mu^-$	[10.09, 12.86]	2.2 ± 0.6	$3.0 \pm 1.5 \pm 1.0$	$2.08^{+0.42}_{-0.39} \pm 0.42$ [11]	$2.17^{+0.57}_{-0.55}$
$\Lambda \mu^+ \mu^-$	[14.18, 16.00]	1.7 ± 0.5	$1.0 \pm 0.7 \pm 0.3$	$2.04^{+0.35}_{-0.33} \pm 0.42$ [11]	1.70 ± 0.44
$\Lambda \mu^+ \mu^-$	> 16.00	7.0 ± 2.9	$7.0 \pm 1.9 \pm 2.2$		7.0 ± 2.9

Results for CDF and LHCb are relative BFs converted to absolute BFs.

† See the original paper for the exact $m^2(\mu^+ \mu^-)$ selection.

‡ The two LHCb measurements include additional binning not reported here.

Heavy FLavor AVeraging group (HFLAV) - May 2018
 Compilation of Ξ_b^0 Branching Fractions ($\times 10^{-6}$)

Preliminary Updated results not included in PDG Live as of Dec. 31, 2017

RPP #	Mode	PDG2017 Avg.	LHCb	Our Avg.
4	$f_{\Xi_b^0}/f_d \mathcal{B}(\Xi_b^0 \rightarrow \bar{K}^0 p \pi^-)$	< 1.6	< 1.6	[1] < 1.6
5	$f_{\Xi_b^0}/f_d \mathcal{B}(\Xi_b^0 \rightarrow \bar{K}^0 p K^-)$	< 1.1	< 1.1	[1] < 1.1
10	$f_{\Xi_b^0}/f_{A_b^0} \mathcal{B}(\Xi_b^0 \rightarrow \Lambda \pi^+ \pi^-)$	< 1.7	< 1.7	[7] < 1.7
11	$f_{\Xi_b^0}/f_{A_b^0} \mathcal{B}(\Xi_b^0 \rightarrow \Lambda K^+ \pi^-)$	< 0.8	< 0.8	[7] < 0.8
12	$f_{\Xi_b^0}/f_{A_b^0} \mathcal{B}(\Xi_b^0 \rightarrow \Lambda K^+ K^-)$	< 0.3	< 0.3	[7] < 0.3
	$f_{\Xi_b^0}/f_{A_b^0} \mathcal{B}(\Xi_b^0 \rightarrow p K^- \pi^+ \pi^-)$	$1.72 \pm 0.21 \pm 0.25 \pm 0.15 \pm 0.07$ [10]	1.72 ± 0.37	
	$f_{\Xi_b^0}/f_{A_b^0} \mathcal{B}(\Xi_b^0 \rightarrow p K^- \pi^+ K^-)$	$1.56 \pm 0.16 \pm 0.19 \pm 0.13 \pm 0.06$ [10]	1.56 ± 0.29	
	$f_{\Xi_b^0}/f_{A_b^0} \mathcal{B}(\Xi_b^0 \rightarrow p K^- K^+ K^-)$	< 0.25	[10] < 0.25	

Channels with no RPP# are not reported by PDG.
 Results for LHCb are relative BFs converted to absolute BFs.

Heavy FLavor AVeraging group (HFLAV) - May 2018
 Compilation of Ξ_b^- Branching Fractions ($\times 10^{-5}$)

Preliminary Updated results not included in PDG Live as of Dec. 31, 2017

RPP #	Mode	PDG2017 Avg.	LHCb	Our Avg.
6	$f_{\Xi_b^-} \mathcal{B}(\Xi_b^- \rightarrow p K^- K^-)/(f_u \mathcal{B}(B^- \rightarrow K^+ K^- K^-))$	†	$265 \pm 35 \pm 47$ [12]	265 ± 58
	$f_{\Xi_b^-} \mathcal{B}(\Xi_b^- \rightarrow p K^- \pi^-)/(f_u \mathcal{B}(B^- \rightarrow K^+ K^- K^-))$		$259 \pm 64 \pm 49$ [12]	259 ± 80
8	$\mathcal{B}(\Xi_b^- \rightarrow p \pi^- \pi^-)/(\mathcal{B}(\Xi_b^- \rightarrow p K^- K^-))$	< 0.56	< 0.56	[12] < 0.56
	$f_{\Xi_b^-} \mathcal{B}(\Xi_b^- \rightarrow p \pi^- \pi^-)/(f_u \mathcal{B}(B^- \rightarrow K^+ K^- K^-))$		< 147	[12] < 147
9	$\mathcal{B}(\Xi_b^- \rightarrow p K^- \pi^-)/(\mathcal{B}(\Xi_b^- \rightarrow p K^- K^-))$	$0.98 \pm 0.27 \pm 0.09$	$0.98 \pm 0.27 \pm 0.09$ [12]	0.98 ± 0.28

Channels with no RPP# are not reported by PDG.
 † PDG reports results multiplied by $\mathcal{B}(B^+ \rightarrow K^+ K^- K^+)$ and $\mathcal{B}(\bar{b} \rightarrow B^+)$.

Heavy FLavor AVeraging group (HFLAV) - May 2018
 Compilation of Ω_b^- Branching Fractions ($\times 10^{-5}$)

Preliminary Updated results not included in PDG Live as of Dec. 31, 2017

RPP #	Mode	PDG2017 Avg.	LHCb	Our Avg.
2	$f_{\Omega_b^-} \mathcal{B}(\Omega_b^- \rightarrow p K^- K^-)/(f_u \mathcal{B}(B^- \rightarrow K^+ K^- K^-))$	†	< 18 [12]	< 18
3	$f_{\Omega_b^-} \mathcal{B}(\Omega_b^- \rightarrow p K^- \pi^-)/(f_u \mathcal{B}(B^- \rightarrow K^+ K^- K^-))$	†	< 51 [12]	< 51
4	$f_{\Omega_b^-} \mathcal{B}(\Omega_b^- \rightarrow p \pi^- \pi^-)/(f_u \mathcal{B}(B^- \rightarrow K^+ K^- K^-))$	†	< 109 [12]	< 109

† PDG reports results multiplied by $\mathcal{B}(B^+ \rightarrow K^+ K^- K^+)$ and $\mathcal{B}(\bar{b} \rightarrow B^+)$.

References

- [1] R. Aaij *et al.*, (LHCb collaboration), JHEP **04**, 087, (2014), arXiv:1402.0770 [hep-ex].
- [2] T. Aaltonen *et al.*, (CDF collaboration), Phys. Rev. Lett. **103**, 031801, (2009), arXiv:0812.4271 [hep-ex].
- [3] T. Aaltonen *et al.*, (CDF collaboration), Phys. Rev. Lett. **107**, 201802, (2011), arXiv:1107.3753 [hep-ex].
- [4] R. Aaij *et al.*, (LHCb collaboration), Phys. Lett. **B725**, 25, (2013), arXiv:1306.2577 [hep-ex].
- [5] D. Acosta *et al.*, (CDF collaboration), Phys. Rev. **D66**, 112002, (2002), arXiv:hep-ex/0208035 [hep-ex].
- [6] R. Aaij *et al.*, (LHCb collaboration), JHEP **09**, 006, (2015), arXiv:1505.03295 [hep-ex].
- [7] R. Aaij *et al.*, (LHCb collaboration), JHEP **05**, 081, (2016), arXiv:1603.00413 [hep-ex].
- [8] R. Aaij *et al.*, (LHCb collaboration), Phys. Lett. **B759**, 282, (2016), arXiv:1603.02870 [hep-ex].
- [9] R. Aaij *et al.*, (LHCb collaboration), JHEP **04**, 029, (2017), arXiv:1701.08705 [hep-ex].
- [10] R. Aaij *et al.*, (LHCb collaboration), JHEP **02**, 098, (2018), arXiv:1711.05490 [hep-ex].
- [11] R. Aaij *et al.*, (LHCb collaboration), JHEP **06**, 115, (2015), arXiv:1503.07138 [hep-ex].
- [12] R. Aaij *et al.*, (LHCb collaboration), Phys. Rev. Lett. **118**, no. 7, 071801, (2017), arXiv:1612.02244 [hep-ex].